

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
22 March 2001 (22.03.2001)

PCT

(10) International Publication Number
WO 01/19158 A2

(51) International Patent Classification: Not classified

(21) International Application Number: PCT/EP00/08875

(22) International Filing Date:
8 September 2000 (08.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
1663/99 10 September 1999 (10.09.1999) CH(71) Applicant (for all designated States except US): CE-
TERIS HOLDING B.V.-AMSTERDAM (OLANDA) -
SUCCURSALE DI LUGANO [CH/CH]; Via Serafino
Balestra, 27, CH-6900 Lugano (CH).

(72) Inventor; and

(75) Inventor/Applicant (for US only): MERIZZI, Gian-
franco [IT/IT]; Via Vela, 7, I-10128 Torino (IT).(74) Agent: RAMBELL, Paolo; Jacobacci & Perani S.p.A.,
Corso Regio Parco, 27, I-10152 Torino (IT).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— Without international search report and to be republished
upon receipt of that report.For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.(54) Title: AN ANTI-OXIDANT PREPARATION BASED ON PLANT EXTRACTS FOR THE TREATMENT OF CIRCULA-
TION AND ADIPOSITOY PROBLEMS(57) Abstract: A preparation based on plant extracts, with an anti-oxidant action which is particularly useful in the prevention
and treatment of circulation problems and in the prevention and treatment of surplus fat deposits, characterised in that its active
ingredients comprise, in association, *Ginkgo bilobabiflavones*, catechine and/or epicatechine, iodine and a component selected from
among madecassic acid, asiatic acid, asiaticoside or combinations thereof.

WO 01/19158 A2

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

(43) International Publication Date
22 March 2001 (22.03.2001)

PCT

(10) International Publication Number
WO 01/19158 A3(51) International Patent Classification?: A61K 35/78.
A61P 9/00

(21) International Application Number: PCT/EP00/08875

(22) International Filing Date:
8 September 2000 (08.09.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
1663/99 10 September 1999 (10.09.1999) CH(71) Applicant (for all designated States except US): CE-
TERIS HOLDING B.V.-AMSTERDAM (OLANDA) -
SUCCURSALE DI LUGANO [CH/CH]; Via Serafino
Balestra, 27, CH-6900 Lugano (CH).(72) Inventor; and
(75) Inventor/Applicant (for US only): MERIZZI, Gian-
franco [IT/IT]; Via Vela, 7, I-10128 Torino (IT).(74) Agent: RAMBELLI, Paolo; Jacobacci & Partners S.p.A.,
Corso Regio Parco, 27, I-10152 Torino (IT).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
21 February 2002For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.(54) Title: AN ANTI-OXIDANT PREPARATION BASED ON PLANT EXTRACTS FOR THE TREATMENT OF CIRCULA-
TION AND ADIPOSITY PROBLEMS(57) Abstract: A preparation based on plant extracts, with an anti-oxidant action which is particularly useful in the prevention
and treatment of circulation problems and in the prevention and treatment of surplus fat deposits, characterised in that its active
ingredients comprise, in association, *Ginkgo biloba* flavonoids, catechine and/or epicatechine, iodine and a component selected from
among modécassic acid, asiatic acid, asiaticoside or combinations thereof.

WO 01/19158 A3

WO 01/19158

PCT/EP00/08875

1

An anti-oxidant preparation based on plant extracts for the treatment of circulation and adiposity problems

The present invention relates to a preparation based on plant extracts which has an anti-oxidant effect and is particularly useful in the treatment of circulation problems such as phlebitis, varicose veins, arteriosclerosis, haemorrhoids and high blood pressure, as well as in the prevention and treatment of surplus fat.

The object of the invention is to provide a preparation to be taken orally, based on a combination of active ingredients of natural and plant origin, which work more effectively to prevent and treat the aforesaid problems when administered by mouth.

This object is achieved according to the invention by providing a preparation characterised in that its active ingredients include a combination of *Ginkgo biloba* biflavones, catechine and/or epicatechine, cumarine and derivatives thereof, iodine and an ingredient chosen from asiaticoside, asiatic acid, madecassic acid and compounds thereof.

The preparation is obtained by mixing plant extracts which contain the above active components.

It is known that extracts from the leaves of *Ginkgo biloba* contain important active components and in particular flavonol glucosides, lactonic terpenes and dimeric biflavones or flavones. The flavonol glucosides and the lactonic terpenes constitute the active components of the standardized *Ginkgo biloba* extracts currently available on the market;

CONFIRMATION COPY

WO 01/19158

PCT/EP00/08875

2

however these extracts do not contain the biflavone component which is not extracted during normal processing. The *Ginkgo biloba* extract used in compositions according to the present invention is highly enriched with the biflavone component and, as a possible option, with extracts containing flavonol glucosides and lactonic terpenes. Five biflavones in particular have been identified in the biflavone component of *Ginkgo biloba*: these are, in particular, amentoflavone, bilobetine, isoginkgetine, ginkgetine and sciadopisidine; the five said compounds differ only by the presence of methyl groups in some positions and, like all flavones, are powerful antioxidants. However, from a pharmacological point of view, they are characterised by their anti-phosphodiesterase, anti-inflammatory, vasculokinetic and anti-allergy properties. Phosphodiesterases (PDE) are cellular enzymes responsible for interacting with cyclic nucleotides so as to linearize them. Cyclic nucleotides are involved as second messengers in transmitting intercellular signals and are thus responsible for some phenomena which are very important from a biochemical point of view. They assist with the visual process and in the relaxation of smooth muscles, they stimulate lipolysis in adiposity and vasculo-motion in capillary arterioles. More specifically, it is sufficient to report that in inhibiting PDE depending on cyclic AMP, these biflavones demonstrate an IC50 of 1.2 micromoles.

The anti-inflammatory properties of biflavones, and in particular those of amentoflavone, have been demonstrated both in vitro, by measuring the interaction of these biflavones with cyclo-oxygenase, lipo-oxygenase and phospholipase A2, and in vivo, using various models of inflammation in animals (carrageneen oedema, Croton oil inflammation etc). The anti-inflammatory action of the biflavones was confirmed

WO 01/19158

PCT/EP00/08875

3

both in models using local application and in those in which they were administered peritoneally. In these models, the biflavones always demonstrated an anti-inflammatory action equivalent to that of indomethacyn or prednisolone. This effectiveness can be explained by analysing the IC50 of cyclo-oxygenase inhibition, which for amentoflavone is 3 micromoles.

With regard to the microvasculokinetic activity of biflavones, it should be reported that, following acute treatment, these substances improve the size of the arterial sphygmie wave and, following chronic treatment they improve capillary density in tissues with trophic-connective problems, such as those affected by panniculopathy and/or various degrees of sclerodermy. Biflavones also have clear anti-allergy properties; they inhibit the release of histamine by mast-cells stimulated by allergens.

In the context of the present invention, it has been demonstrated that, when administered orally, the activity of the aforesaid biflavones, possibly in combination with flavonol glucosides and lactonic terpenes which are normally present in standard *Ginkgo biloba* extracts, is enhanced when the latter are combined with the aforesaid active compounds.

The extracts are preferably used in a phytosomal form, in which the active components are compounded with phospholipids.

In the context of the invention it is convenient to use an extract of leucocyanidine or leucoanthocyanine derived from *Vitis vinifera* as the source of catechine or epicatechine. Leucoanthocyanines are procyanidolic oligomers derived from

WO 01/19158

PCT/EP00/08875

4

condensing monomeric units of flavan-3-ols and flavan-3,4-diols, these being either free or esterified with gallic acid. Leucocyanines are powerful anti-oxidant substances with the ability to protect capillaries by increasing oxygen to the tissue; these active substances prove biologically active even when administered orally and they have been shown to be tropic for the cardio-vascular system and for all tissues, such as artery walls, which are rich in glucoaminoglycene. Preferably, phytosomal forms of extracts are used, thus further enhancing the bioavailability of the active principles. In this form the procyanodines are completed with phospholipids, particularly distearylphosphatidylcholine of soya.

The preferable source of cumarine is an extract of *Melilotus* (*Melilotus officinalis*), cumarine and its derivatives being the main active ingredients thereof; the main active ingredients of this extract are melilotine (3,4 dihydro-cumarine), melilotic acid (hydroxycumarinic acid), melilotoxide (a melilotine glucoside) and some flavinoids which act like vitamin P; the active ingredients contained in the extract are particularly effective in increasing capillary strength, in reducing vascular permeability, in stimulating venous circulation and improving lymphatic circulation.

Extract of *Melilotus* may be replaced or backed up, as a source of cumarine and its derivatives, by an extract of *Aesculus hippocastanum* (horse chestnut) in the same dosage or up to around twice the dose of *Melilotus* extract.

The most abundant active principle of *Aesculus hippocastanum* extract, obtained from the bark, the pericarp of the fruit,

WO 01/19158

PCT/EP00/08875

5

the leaves or the buds, is coumarine glucoside, esculoside (6-0-glucosil-7-hydroxy-coumarine). Other coumarines contained in the extract are fraxine (8-0-glycoside-7-hydroxy-6-metoxycoumarine) and the aglicones, esculetin (6,7-dioxy-coumarine) and fraxetine (7,8-dioxy-6-methoxy-coumarine).

The preferred source of asiaticoside, asiatic acid and madecassic acid is an extract containing a triterpenic fraction of centella (*Centella Asiatica*) which contains a combination of the above three active principles. The extract should preferably be used in a phytosomal form, obtained by a reaction between the triterpenic fraction of *Centella Asiatica* with a phospholipid. A main action of the triterpenic fraction of centella consists in accelerating the uptake and metabolism of lysine and of proline, thus increasing the synthesis and the release of tropocollagen and stimulating the turnover of acid mucopolysaccharide acids in connective tissue. Thanks to these properties, the active principles are particularly effective in reducing localised adiposity.

The preferred source of iodine is an extract of *Fucus vesiculosus*, a seaweed of the Fucaceae family. The role of the iodine in *Fucus vesiculosus* is currently well characterised; it is made more readily available by complexing with a protein fraction of the extract; it increases the synthesis of thyroid hormones and indirectly enhances the lipolytic action of these hormones (T3 and T4) thanks to a local thermogenic action on adipose tissue. The *Fucus* extract also contains polysaccharides such as fucoidin, alginic acid, laminarin and poliose; of these, the alginic acids, in particular, enhance the effect of the extract since they are hydrophilic molecules able to swell from their

WO 01/19158

PCT/EP00/08875

6

original volume in the presence of water; thanks to this characteristic, when ingested they not only reduce the appetite but also increase the speed of transit of food through the intestines, thus ensuring that it is less well absorbed.

The basic composition of the invention can thus be obtained by mixing a biflavone extract of *Ginkgo biloba* (perhaps in combination with a standard *Ginkgo biloba* extract also containing flavonol glucosides and lactonic terpenes), leucocyanidine extract, Melilotus extract, Centella extract and extract of *Fucus vesiculosus*; these extracts preferably being in a phytosome form.

With reference to the extracts normally available on the market, the basic composition is preferably made up by the following percentages by weight:

- 1.5 - 32% biflavone extract of *Ginkgo biloba*;
- 6 - 80% of leucocyanidine extract;
- 1.5 - 60%, preferably 1.5 - 32% of melilotus extract and/or *Aesculus hippocastanum* extract;
- 1.5 - 32% of centella extract;
- 1.5 - 85% *Fucus vesiculosus* extract, possibly in combination with:
- 1.5 - 32% of standard *Ginkgo biloba* extract containing flavon glucosides and lactonic terpenes.

In terms of the content of active principles, the composition of the invention preferably contains the following percentages by weight:

- 0.2 - 12%, preferably 0.5 - 6% of total biflavones, expressed as ginkgetine content;

WO 01/19158

PCT/EP00/08875

7

0.2 - 12%, preferably 0.5 - 8% of catechine and/or epicatechine, expressed as catechine content;

0.15 - 7%, preferably 0.3 - 3.5% of cumarine and its derivatives;

0.25 - 15%, preferably 0.5 - 7.5% of asiaticoside;

0.25 - 22%, preferably 0.5 - 11 % of asiatic acid and/or madecassic acid;

up to 0.5% of iodine and possibly one or more of its components:

0.25 - 12%, preferably 0.5 - 6%, of flavonol glucosides, and

0.1 - 2%, preferably 0.2 - 1% of Ginkgolide lactonic terpenes (bilobalide).

The composition can also contain active ingredients chosen from methylxanthinic derivatives, such as caffeine, theobromine or theophylline in particular; mono and di-caffeoylchinic acids, chlorogenic acids, eicosapentaenoic acid (EPA), docahexaenoic acid (DHA) gamma-linolenic acid and combinations thereof.

The preferred source of methylxanthinic derivatives, of caffeine and of chlorogenic acids in particular, is an extract of *Ilex paraguariensis*, possibly in phytosomal form; the standard, commercially available extract can be added to the previously-described basic mixture in a quantity of 1.5 to 32% by weight.

Caffeoylchinic acid and its derivatives (such as cynarine) are preferably introduced by means of an artichoke extract (*Cynara scolymus*); this extract typically also contains caffeic acid and chlorogenic acid; this extract is preferably

WO 01/19158

PCT/EP00/08875

8

used in quantities of 1.5 to 32% by weight with reference to 100 parts of the basic mixture.

The preferable source of eicosapentaenoic acid (EPA) and of docohexaenoic acid (DHA) is fish oil which, with reference to 100 parts of the basic mixture, may be added in quantities of 5 to 80% by weight.

Gamma-linolenic acid is preferably introduced into the formulation by the use of borage oil, added in quantities of 30 to 120% by weight with reference to 100 parts of basic preparation.

In particular, in the preferred embodiment of the invention, the composition includes one or more of the following components:

Caffeine in quantities of 0.05 to 2.5 (preferably 0.1 - 1%) by weight with reference to the total preparation,

Caffeoylchinnic acids in quantities from 0.05 to 4.8% (preferably 0.01 to 2.5%) by weight,

Eicosapentanoic acid in quantities from 0.75 to 24% (preferably from 1.5 to 12%) by weight,

Docohexanoic acid in quantities from 0.6 to 8% (preferably 1.2 to 4%) by weight,

Gamma-linolenic acid in quantities from 2.5 to 22%, preferably 5-11% by weight.

For example, a typical composition could be formulated according to the data in the table below, which gives the preferred minimum and maximum quantities expressed in parts by weight of the components of the basic mixture (marked with an asterisk) and of optional ingredients.

WO 01/19158

PCT/EP00/08875

9

| | Minimum (Parts by weight) | Maximum (Parts by weight) |
|--|---------------------------------|---------------------------------|
| *Dry extract of <i>Vitis vinifera</i> (optionally phytosomes) | 20 | 200 |
| *Dry extract of <i>Melilotus officinalis</i> and/or <i>Aesculus hippocastanum</i> | 5 | 40 |
| * <i>Ginkgo biloba</i> biflavones (optionally phytosomes) | 5 | 50 |
| Dry extract of <i>Ginkgo biloba</i> (optionally phytosomes) | 5 | 50 |
| *Dry extract of <i>Fucus vesiculosus</i> | 50 | 200 |
| *Dry extract of <i>Centella asiatica</i> (optionally phytosomes) | 10 | 50 |
| Dry extract of artichoke | 10 | 100 |
| Dry extract of <i>Ilex paraguariensis</i> | 10 | 100 |
| Borage oil | 50 | 1000 |
| Fish oil | 50 | 750 |
| Soya lecithin | 20 | 100 |

In the above table, the given values, expressed in parts by weight, correspond, when expressed in milligrams to the minimum and maximum daily doses recommended or to the dose per capsule.

WO 01/19158

PCT/EP00/08875

10

The composition of the invention is formulated in forms suited to be taken orally, such as, for example, gelatin capsules with either soft or rigid shells, tablets, pills, elixirs, suspensions and syrups. The various forms can include excipients and/or binders and/or pharmaceutically acceptable vehicles, in particular lecithin mono- and diglycerides of fatty acids. The mixture of extracts can be administered orally, possibly in an edible vehicle or can be incorporated directly into food as part of a diet.

The preparation is particularly useful in treating localised adiposity, in particular in men but also in women. It is well known that adiposity differs in men and in women, by the area in which it is deposited, by the quantity, by functional response and by consequences for health; in actual fact this surplus fat ("spare tyres" in men and cellulite in women) is due, at least in part, to a tropism which is known to involve both vascular and connective tissue, as well as adipose tissue. The composition of the invention provides an association of well characterised substances from a pharmacotoxicological and chemical point of view, which is totally free of side effects and is particularly advantageous compared to pharmaceutical preparations based on appetite-suppressant compounds, as used in conventional pharmacological treatment.

WO 01/19158

PCT/EP00/06875

11

CLAIMS

1. A composition based on plant extracts, with an anti-oxidant activity which is particularly useful in the prevention and treatment of circulation problems and in the prevention and treatment of adiposity, characterised in that its active ingredients comprise, in association, biflavones of *Ginkgo biloba*, catechine and/or epicatechine, cumarine or derivatives thereof, iodine and a component chosen from among madecassic acid, asiatic acid, asiaticoside and mixtures of these.
2. A composition according to Claim 1, characterised in that it is obtained by mixing plant extracts containing the aforesaid active principles.
3. A composition according to Claim 2, characterised in that the said extracts are in phytosomal form.
4. A composition according to any one of the preceding Claims, characterised in that it further includes flavonol glucosides and lactonic terpenes.
5. A composition according to any Claim from 1 to 4, characterised in that it also includes an active principle chosen from the group consisting of methylxanthinic derivatives, chlorogenic acids, mono- and di-caffeoylchinnic acid and derivatives thereof, eicosapentaenoic acid, docohexaenoic acid, gamma-linolenic acid and mixtures thereof.

WO 01/19158

PCT/EP00/08875

12

6. A composition according to any of Claims 1 to 4, characterised in that it is obtained by mixing plant extracts in the following percentages by weight:

1.5-32% of *Ginkgo biloba* biflavone extract;
6-80% of leucocyanidine;
1.5-32% of *Melilotus* and/or *Aesculus*
hyppocastanum extract;
1.5-32% of centella extract;
1.5-85% extract of *Fucus vesiculosus*; and
optionally
1.5-32% of standardised *Ginkgo biloba* extract
containing flavone glucosides and lactonic terpenes.

7. A composition according to Claim 6, characterised in that with reference to 100 parts of the basic mixture of Claim 6, it also includes one or more of the following components:

from 1.5 to 32% by weight of *Ilex paraguariensis* extract;

from 1.5 to 32% by weight of artichoke extract;

from 5 to 80% by weight of fish oil;

and

from 30 to 120% by weight of borage oil.

8. A composition according to any one of the preceding Claims which includes:

0.12-12%, preferably 0.5-6% by weight, of total biflavones;

0.2-12%, preferably 0.5-8% by weight, of catechine and/or epicatechine;

0.15-7%, preferably 0.3-3.5% by weight, of coumarins and derivatives thereof;

0.25-15%, preferably 0.5-7.5% by weight of asiaticoside;

WO 01/19158

13

CT/EP00/08875

0.25-22%, preferably 0.5-11% by weight, of asiatic acid and/or madecassic acid;

up to 0.5% of iodine and, optionally

0.25-12%, preferably 0.5-6% by weight, of flavonol glucosides and

0.1-2%, preferably 0.2-1% by weight, of lactonic terpenes.

9. A composition according to Claim 8, characterised in that it also includes one or more of the following components:

from 0.05 to 2.5%, preferably 0.1-1% by weight, of caffeine;

from 0.05 to 4.8%, preferably 0.1-2.5% by weight, of caffeilchinic acids;

from 0.75 to 24%, preferably 1.5-12% by weight, of eicosapentaenoic acid;

from 0.6 to 8%, preferably 1.2-4% by weight, of docoesaenoic acid; and

from 2.5 to 22%, preferably from 5 to 11% by weight of gamma-linolenic acid.

10. A composition according to any one of the preceding Claims in a pharmaceutical form for oral administration.

11. The use of flavone dimers in the formulation of a preparation based on plant extracts useful in the prevention and treatment of circulation problems and adiposity.

INTERNATIONAL SEARCH REPORT

Int'l. Classification No.

PCT/EP 00/08875

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K35/78 A61P9/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data, BIOSIS, CHEM ABS Data, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No |
|------------|---|----------------------|
| P, X | W. REILLY, V. REEVE: "Body contouring using an oral herbal antioxidant formulation-Centelaplus: a dose controlled observational study" REDOX REPORT, vol. 5, no. 2-3, 2000, pages 144-145, XP000990069 page 144 | 1, 2, 5, 10, 11 |
| A | GB 2 174 904 A (SEUREF) 19 November 1986 (1986-11-19) claims 1, 3, 4, 6 | 1 |

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *Z* document member of the same patent family

Date of the actual completion of the international search

11 June 2001

Date of mailing of the international search report

22/06/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Peeters, J

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Application No.

PCT/EP 80/08875

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| GB 2174904 A | 19-11-1986 | CH 662505 A | 15-10-1987 |
| | | AT 393793 B | 10-12-1991 |
| | | AT 115186 A | 15-06-1991 |
| | | BE 904696 A | 18-08-1986 |
| | | DE 3614278 A | 30-10-1986 |
| | | FR 2587212 A | 20-03-1987 |
| | | JP 1679881 C | 13-07-1992 |
| | | JP 3044048 B | 04-07-1991 |
| | | JP 61280424 A | 11-12-1986 |
| | | NL 8601105 A | 17-11-1986 |
| | | US 4707360 A | 17-11-1987 |